

Predictability and Dynamics of Geophysical Fluid Flows

Roger M. Samelson

College of Oceanic and Atmospheric Sciences, Oregon State University

104 Ocean Admin Bldg

Corvallis, OR 97331-5503 USA

phone: (541) 737-4752 fax: (541) 737-2064 email: rsamelson@coas.oregonstate.edu

Award Number: N00014-98-1-0813

<http://www.coas.oregonstate.edu/~faculty/samelson.html>

LONG-TERM GOALS

The long-term goal of my research in this project is to improve our ability to predict environmental conditions using dynamical models.

OBJECTIVES

The central objective of my research in this project is to understand the mathematical and physical connections between the bred-growing-mode and singular vector techniques recently developed for numerical weather prediction, the Lyapunov vectors and exponents of dynamical systems theory, and instability theories of geophysical fluid dynamics. My intent is to gain insight into fundamental mathematical and physical aspects of predictability in unstable (irregular, chaotic) continuous systems.

APPROACH

I am using a combination of analytical and numerical methods to study a variety of mathematical models of geophysical fluid flows.

WORK COMPLETED

Graduate student Christopher Wolfe, who is supported by this grant, and I have completed and published a study of a strongly nonlinear baroclinic wave-mean oscillation and its time-dependent normal-mode instabilities in a high-dimensional geophysical fluid model (Samelson and Wolfe, 2003). This model is a two-layer, quasi-geostrophic, numerical channel model with 48 along-channel and 40 cross-channel modes in each layer, for a total of 3,840 degrees of freedom. This study included a novel application of the efficient Newton-Picard solver PDECONT developed by Lust et al. (1998). We have also completed the study of a number of related nonlinear wave-mean cycles, for which we have computed all of the unstable and stable linear disturbance eigenvalues and eigenmodes.

This project has also provided partial support for several other efforts, including research on Lagrangian motion in coastal ocean circulation (Kuebel et al., 2003; Kuebel Cervantes et al., 2003), and studies of the coastal lower atmosphere and comparisons of model and scatterometer wind stress fields (Perlin et al., submitted). Wolfe was a Student Fellow at the 2003 Woods Hole Summer Program in Geophysical Fluid Dynamics, where he worked with C. Cenedese on laboratory models of eddy generation by flow over variable topography (Wolfe, 2003).

Report Documentation Page				Form Approved OMB No. 0704-0188	
Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.					
1. REPORT DATE 30 SEP 2003		2. REPORT TYPE		3. DATES COVERED 00-00-2003 to 00-00-2003	
4. TITLE AND SUBTITLE Predictability and Dynamics of Geophysical Fluid Flows				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S)				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) College of Oceanic and Atmospheric Sciences, Oregon State University,,104 Ocean Admin Bldg,,Corvallis,,OR,97331				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release; distribution unlimited					
13. SUPPLEMENTARY NOTES					
14. ABSTRACT					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT Same as Report (SAR)	18. NUMBER OF PAGES 6	19a. NAME OF RESPONSIBLE PERSON
a. REPORT unclassified	b. ABSTRACT unclassified	c. THIS PAGE unclassified			

RESULTS

The complete Floquet spectra computed for strongly nonlinear cycles show structure similar to that found in the weakly nonlinear case (Samelson, 2001): a neutral wave-dynamical mode (Figure 1), several weakly damped or unstable wave-dynamical modes (Figure 2), several rapidly damped wave-dynamical modes (Figure 3), and many modes that are damped at a rate roughly equal to the Ekman friction timescale. The latter are often dominated by small-scale features that are advected along the channel in opposite directions by the mean flow in each layer (Figure 4). This behavior is in marked contrast to the wave-dynamical modes, which are coherent between the layers and are quasi-stationary in these flows with nearly zero mean barotropic motion. Inclusion of higher-order viscosity induces additional scale selectivity in the frictionally damped modes.

IMPACT/APPLICATIONS

The primary potential future impact of these results is on the design and use of ensemble forecasting techniques for the prediction of oceanic and atmospheric conditions.

RELATED PROJECTS

This work formed part of the ONR Predictability DRI. The coastal meteorological research is partially supported by the ONR project “COAMPS Simulations of the Coastal Atmosphere” and the NSF project “COAST: Coastal Ocean Advances in Shelf Transport.” The research on dynamics and predictability is related to the NSF ITR project “Spatio-Temporal Complexity and Nonlinear Dynamics of Coastal Ocean Flows” (R. Samelson, J. Allen, and G. Egbert, Co-PIs).

SUMMARY

The results described above open a new perspective on the analysis of the evolution and predictability of oceanic and atmospheric flows, by showing that techniques previously restricted to highly simplified models can be extended and adapted for models that are sufficiently complex that they can be expected to provide substantial insight into geophysical fluid motion. This perspective is yielding new results relevant to instability theory and to ensemble forecasting methods for environmental prediction. Participation in this ONR-sponsored work has enhanced my institution by increasing its visibility in the research community, stimulating interaction among institutional colleagues, and supporting a graduate student specifically interested in the problems addressed in this work.




Figure 1. Contours (red-positive; blue-negative; absolute amplitude arbitrary) of streamfunction in layers 1 (upper panel) and 2 (lower panel) along the center of the periodic zonal channel vs. time (one oscillation period) for neutral linear mode of a nonlinear wave-mean oscillation




Figure 2. As in Fig. 1, but for the least stable linear mode of the nonlinear wave-mean oscillation. The mode is normalized by its exponential growth factor, so the contours show the time-periodic structure function.




Figure 3. As in Fig. 2, but for the corresponding rapidly-damped wave-dynamical linear mode of the nonlinear wave-mean oscillation. Note the approximate time-reversal symmetry relative to the mode in Fig. 2.




Figure 4. As in Fig. 2, but for an advective, frictionally damped linear mode of the nonlinear wave-mean oscillation. The mean flow is to the right in layer 1 (upper panel) and to the left in layer 2 (lower panel).

REFERENCES

Lust, K., D. Roose, A. Spence, and A. Champneys, 1998. An adaptive Newton-Picard algorithm with subspace iteration for computing periodic solutions. *SIAM Journal on Scientific Computing*, 19, 1188-1209.

PUBLICATIONS

Kuebel, B. T., J. S. Allen, P. A. Newberger, and R. M. Samelson, 2003. A modeling study of Eulerian and Lagrangian aspects of shelf circulation off Duck, North Carolina. *Journal of Physical Oceanography*. [in press]

Kuebel Cervantes, B. T., J. S. Allen, and R. M. Samelson, 2003. Lagrangian characteristics of continental shelf flows forced by periodic wind stress. *Nonlinear Processes in Geophysics*, submitted.

Perlin, N., R. M. Samelson, and D. B. Chelton, Scatterometer and model wind and wind stress in the Oregon-California coastal zone. *Monthly Weather Review*, submitted.

Samelson, R. M., and C. L. Wolfe, 2003. A nonlinear baroclinic wave-mean oscillation with multiple normal-mode instabilities. *Journal of the Atmospheric Sciences*, 60, 1186-1199. [published]

Wolfe, C. L., 2003. Eddy generation by flow over variable topography: some experiments. In *Proceedings of 2003 Summer Program in Geophysical Fluid Dynamics*, Woods Hole Tech. Rep., to appear. [in press]

AWARDS

Christopher L. Wolfe, Student Fellowship, Woods Hole Summer Program in Geophysical Fluid Dynamics